

# Curvature and the Electromagnetic Field

Craig W. Brown

13512 Wallingford Avenue N, Seattle, WA 98133

*Corresponding author E-mail:* cbphysics@comcast.net

(Received 20 March 2021, Accepted 11 April 2021, Published 29 April 2021)

## Abstract

Maxwell's equations are derived from the curvature tensor and a vector potential. The results are combined with Einstein's equations. Complete solutions to the resulting equation yield simultaneous solutions to both Einstein's and Maxwell's equations. This is a classical theoretical unification of electromagnetism and gravitation.

**Keywords:** Electromagnetism; Curvature Tensor; Maxwell's Equations; General Relativity; Unified Field Theory

Since the geometrical general theory of relativity was published as a theory of gravitation [1], the notion that electromagnetism was also amenable to a classical, geometrical theory has been pursued in a variety of ways [2], including notable contributions by Weyl [3], Kaluza, and Schrödinger [4]. To date, however, the achievement of this goal is not generally accepted. This is largely due to the fact that the metric tensor is symmetric while the electromagnetic field tensor is antisymmetric. This paper presents a geometrical theory of electromagnetism that is distinct from the various approaches developed by others and introduces the required antisymmetry in a straightforward way. The result is that Maxwell's equations follow, thus unifying the core of classical electromagnetic theory with Einstein's classical gravitational theory. The sole purpose of the present paper is to present this derivation.

The present geometrical theory of electromagnetism begins with the conventional derivation of the fourth-rank curvature tensor  $R_{\nu\rho}^{\beta}$  in four-dimensional spacetime [5]. The lower-case Greek suffixes take on the values 0,1,2,3; downstairs suffixes denote covariant character; upstairs suffixes denote contravariant character. The summation convention for contraction over pairs of covariant and contravariant components with the same Greek suffix is used.

This derivation begins with a covariant vector  $A_\nu$ . The first and second covariant derivatives of the vector are  $A_{\nu:\rho:\sigma}$  where the colon symbol represents covariant differentiation. Using the symbol  $\langle\rho\sigma\rangle$  to denote the tensor preceding it with the suffixes  $\rho$  and  $\sigma$  interchanged, the curvature tensor is defined in the following expression:

$$A_{\nu:\rho:\sigma} - \langle\sigma\rho\rangle = A_{\nu:\rho:\sigma} - A_{\nu:\sigma:\rho} = A_\beta R_{\nu\rho\sigma}^\beta.$$

Only the interchange of  $\rho$  with  $\sigma$  is used above.

The derivation then employs a sum of the cyclic permutations (c.p.) of the three suffixes in the above equation

$$\begin{aligned} A_{\nu:\rho:\sigma} - A_{\nu:\sigma:\rho} + c.p. &= A_{\nu:\rho:\sigma} - A_{\nu:\sigma:\rho} + A_{\rho:\sigma:\nu} - A_{\sigma:\rho:\nu} + A_{\sigma:\nu:\rho} - A_{\rho:\nu:\sigma} \\ &= A_\beta (R_{\nu\rho\sigma}^\beta + R_{\rho\sigma\nu}^\beta + R_{\sigma\nu\rho}^\beta). \end{aligned} \quad (0.1)$$

With inherent symmetries in the curvature tensor,

$$R_{\nu\rho\sigma}^\beta + R_{\rho\sigma\nu}^\beta + R_{\sigma\nu\rho}^\beta = 0$$

and the right-hand side of Eq. (0.1) equals zero [6].

The form of the left-hand side of Eq. (0.1) is simplified using the antisymmetric tensor formed from the covariant derivatives of the vector,  $B_{\nu\rho} = A_{\nu:\rho} - A_{\rho:\nu}$ . With this tensor, the various second derivatives in Eq. (0.1) can be grouped pair-wise based on the suffixes of the second derivatives to become

$$B_{\nu\rho:\sigma} + B_{\rho\sigma:\nu} + B_{\sigma\nu:\rho} = B_{\nu\rho:\sigma} + c.p. = 0. \quad (0.2)$$

This result is a consequence solely of the spacetime geometry. The vector  $A_\nu$  can be any vector and the result is independent of any physical units which the vector may have. If the metric tensor is dimensionless and the vector  $A_\nu$  has the units of the electromagnetic vector potential, which are *tesla·meter* in SI units [7], then the antisymmetric tensor  $B_{\nu\sigma}$  has units of the electromagnetic field and Eq. (0.2) gives two of Maxwell's equations [8], [9].

The antisymmetric tensor formed from the covariant derivatives of a vector is the same as the antisymmetric tensor formed from the ordinary derivatives of the vector,

$$A_{\nu:\rho} - A_{\rho:\nu} = A_{\nu,\rho} - \Gamma_{\nu\rho}^\alpha A_\alpha - A_{\rho,\nu} + \Gamma_{\rho\nu}^\alpha A_\alpha = A_{\nu,\rho} - A_{\rho,\nu}$$

where a comma denotes ordinary differentiation and the  $\Gamma_{\nu\rho}^\alpha$  are Christoffel symbols of the second kind. In similar fashion, it is interesting to write out Eq. (0.2) as follows:

$$\begin{aligned} B_{\nu\rho:\sigma} + c.p. &= \left( A_{\nu,\rho} - \Gamma_{\nu\rho}^\alpha A_\alpha - A_{\rho,\nu} + \Gamma_{\rho\nu}^\alpha A_\alpha \right)_{;\sigma} + c.p. \\ &= B_{\nu\rho,\sigma} - \Gamma_{\nu\sigma}^\beta B_{\beta\rho} - \Gamma_{\sigma\rho}^\beta B_{\nu\beta} + B_{\rho\sigma,\nu} - \Gamma_{\rho\nu}^\beta B_{\beta\sigma} - \Gamma_{\nu\sigma}^\beta B_{\rho\beta} + B_{\sigma\nu,\rho} - \Gamma_{\sigma\rho}^\beta B_{\beta\nu} - \Gamma_{\rho\nu}^\beta B_{\sigma\beta} \\ &= B_{\nu\rho,\sigma} + B_{\rho\sigma,\nu} + B_{\sigma\nu,\rho} - \Gamma_{\nu\sigma}^\beta (B_{\beta\rho} + B_{\rho\beta}) - \Gamma_{\sigma\rho}^\beta (B_{\nu\beta} + B_{\beta\nu}) - \Gamma_{\rho\nu}^\beta (B_{\beta\sigma} + B_{\sigma\beta}) \\ &= B_{\nu\rho,\sigma} + c.p. = 0. \end{aligned}$$

Thus, the gravitational field embodied in the Christoffel symbols does not enter into the Maxwell's equations of Eq. (0.2).

The remaining Maxwell's equations arise following contraction over one suffix in the antisymmetric tensor with the suffix of the covariant differentiation in  $B_{\nu\rho:\sigma}$  using the contravariant metric tensor  $g^{\rho\sigma}$ :

$$g^{\rho\sigma} B_{\nu\rho:\sigma} = \mu_0 J_\nu, \quad (0.3)$$

where  $\mu_0$  is the permeability of a vacuum [9], [10].

The electromagnetic current  $J_\nu$  is then expressible in terms of the vector potential and the Ricci tensor [11]:

$$g^{\rho\sigma} B_{\nu\rho:\sigma} = A_{\nu:\rho}^{:\rho} - g^{\rho\sigma} (A^\beta R_{\beta\rho\nu\sigma} + A_{\rho:\sigma:\nu}^{:\rho}) = \square A_\nu + A^\beta R_{\beta\nu} - (A_\rho^{:\rho})_{;\nu} = \mu_0 J_\nu,$$

where  $\square$  is the covariant D'Alembertian operator such that  $A_{\nu:\rho}^{:\rho} = \square A_\nu$ . This result is general for the vector potential  $A_\nu$  in any gauge. The result is simplified in the covariant Lorentz gauge for which  $A_\rho^{:\rho} = 0$ :

$$\mu_0 J_\nu = \square A_\nu + A^\beta R_{\beta\nu}. \quad (0.4)$$

Unlike the first two Maxwell's equations, the source term equations of Eq. (0.3) and Eq. (0.4) do depend on the curvature. There is curvature at a point if there is any matter at that point, matter being any matter-energy aside from the gravitational field. In the case where sources are present at a point, the curvature obtains contributions due not only from the electromagnetic field but also from the masses of the charged particles of the current. In the case where no sources are present, no other matter is present, and only the free electromagnetic field is present, the curvature obtains contributions solely from the free electromagnetic field and Eq. (0.4) becomes

$$0 = \square A_\nu + A^\beta R_{\beta\nu}.$$

These results can be applied in Einstein's equations for the gravitational field (neglecting the cosmological constant),

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = \kappa T_{\mu\nu} \quad (0.5)$$

Here  $g_{\mu\nu}$  is the metric tensor,  $R$  is the scalar curvature  $g^{\alpha\beta} R_{\alpha\beta}$ ,  $T_{\mu\nu}$  is the matter-energy tensor, and  $\kappa$  is the Einstein gravitational constant that has units of inverse force and equals  $8\pi G/c^4 \approx 2.08 \times 10^{-43} N^{-1}$ , where  $G$  is Newton's constant of gravitation and  $c$  is the speed of light in a vacuum [11]. Using Eq. (0.4), the inner product of a vector potential  $A^\mu$  with Eq. (0.5) yields

$$\mu_0 J_\nu - \square A_\nu - \frac{1}{2} R A_\nu = \kappa A^\mu T_{\mu\nu}.$$

This equation can be solved for the vector potential,

$$A_\nu = \frac{2(\mu_0 J_\nu - \square A_\nu - \kappa A^\mu T_{\mu\nu})}{R}.$$

This result cannot become singular when the scalar curvature is zero. This is because this equation is applicable only when there is matter at least in the form of an electromagnetic field; the curvature cannot then be zero.

At a point where the only matter-energy is the free electromagnetic field, the matter-energy tensor is the electromagnetic stress-energy tensor [12],

$$T_{\mu\nu} = \frac{1}{\mu_0} \left( B_{\mu\alpha} B_\nu{}^\alpha - \frac{1}{4} g_{\mu\nu} B_{\alpha\beta} B^{\alpha\beta} \right).$$

This stress-energy tensor depends only upon the metric, the permeability constant, and derivatives of the vector potential. Given the constants, Einstein's equations for the gravitational field now yield a differential equation for the vector of the free electromagnetic field,

$$\frac{2 \left\{ \square A_\nu + \frac{\kappa}{\mu_0} A^\mu \left[ (A_{\mu:\alpha} - A_{\alpha:\mu}) (A_\nu^{:\alpha} - A^{\alpha:\nu}) - \frac{1}{4} g_{\mu\nu} (A_{\alpha:\beta} - A_{\beta:\alpha}) (A^{\alpha:\beta} - A^{\beta:\alpha}) \right] \right\}}{R} + A_\nu = 0.$$

This equation depends, also, on the metric tensor. Thus, complete solutions to this equation yield simultaneous solutions to both Einstein's and Maxwell's equations for the free electromagnetic field. At points where the current is  $J_\nu \neq 0$ , the differential equation becomes more complicated and may depend upon additional constants, such as mass, and particle velocities associated with the current at such points that may contribute to the matter-energy tensor.

The results presented in this paper employ only the well-established geometric interpretation of general relativity and the conventional use of a vector potential for the electromagnetic field.

## References

- [1] Albert Einstein, *Annalen der Physik*, **49**, (1916); translation by W. Perrett and G. B Jeffery, *The Principle of Relativity*, (Dover Publications, New York, 1952) pp. 111-164.
- [2] Wikipedia contributors. (2020, December 13). Classical unified field theories. In *Wikipedia, The Free Encyclopedia*. Retrieved 00:57, March 3, 2021, ([https://en.wikipedia.org/w/index.php?title=Classical\\_unified\\_field\\_theories&oldid=993884054](https://en.wikipedia.org/w/index.php?title=Classical_unified_field_theories&oldid=993884054))
- [3] Hermann Weyl, *Space-Time-Matter* (translation by Henry L. Brose, Dover Publications, Inc., New York, 1952).
- [4] Erwin Schrödinger, *Space-Time Structure* (Cambridge University Press, 1950, reprinted 1994).
- [5] Paul Adrien Maurice Dirac, *General Theory of Relativity* (Wiley, New York, 1975), Chap. 11, p20-21.
- [6] Paul Adrien Maurice Dirac, *General Theory of Relativity* (Wiley, New York, 1975), Eq. (11.5), p21.
- [7] Richard E. Moss, *Advanced Molecular Quantum Mechanics* (Chapman and Hall, London, 1973), Appendix A, p265-269.
- [8] Paul Adrien Maurice Dirac, *General Theory of Relativity* (Wiley, New York, 1975), Eq. (23.12), p43.
- [9] Richard E. Moss, *Advanced Molecular Quantum Mechanics* (Chapman and Hall, London, 1973) Eq. (5.55), p75.
- [10] Paul Adrien Maurice Dirac, *General Theory of Relativity* (Wiley, New York, 1975), Eq. (23.13), p43.
- [11] Wikipedia contributors. (2021, March 6). Einstein field equations. In *Wikipedia, The Free*

- Encyclopedia*. Retrieved 15:53, March 20, 2021,  
[\(\[https://en.wikipedia.org/w/index.php?title=Einstein\\\_field\\\_equations&oldid=1010723186\]\(https://en.wikipedia.org/w/index.php?title=Einstein\_field\_equations&oldid=1010723186\)\)](https://en.wikipedia.org/w/index.php?title=Einstein_field_equations&oldid=1010723186)
- [12] Wikipedia contributors. (2021, February 11). Electromagnetic stress–energy tensor.  
In *Wikipedia, The Free Encyclopedia*. Retrieved 15:39, March 20, 2021,  
[\(\[https://en.wikipedia.org/w/index.php?title=Electromagnetic\\\_stress%20%93energy\\\_tensor&oldid=1006108690\]\(https://en.wikipedia.org/w/index.php?title=Electromagnetic\_stress%20%93energy\_tensor&oldid=1006108690\)\)](https://en.wikipedia.org/w/index.php?title=Electromagnetic_stress%20%93energy_tensor&oldid=1006108690)